arXiv:1608.05798v1 [math.AG] 20 Aug 2016

NATURALITY OF THE HYPERHOLOMORPHIC SHEAF OVER THE CARTESIAN SQUARE OF A MANIFOLD OF $K3^{[n]}$-TYPE

EYAL MARKMAN

Abstract. Let $\mathcal{M}$ be a $2n$-dimensional smooth and compact moduli space of stable sheaves on a $K3$ surface S and $\mathcal{U}$ a universal sheaf over $S \times \mathcal{M}$. Over $\mathcal{M} \times \mathcal{M}$ there exists a natural reflexive sheaf E of rank $2n-2$, namely the first relative extension sheaf of the two pullbacks of $\mathcal{U}$ to $\mathcal{M} \times S \times \mathcal{M}$. We prove that E is $\omega \boxplus \omega$-slope-stable with respect to every Kähler class ω on $\mathcal{M}$. The sheaf E is known to deform to a sheaf E' over $X \times X$, for every manifold X deformation equivalent to $\mathcal{M}$, and we prove that E' is $\omega \boxplus \omega$-slope-stable with respect to every Kähler class ω on X. This triviality of the stability chamber structure combines with a result of S. Mehrotra and the author to show that the deformed sheaf E' is canonical; each component of the the moduli space of marked triples (X, η, E'), where η is a marking of $H^2(X, \mathbb{Z})$, maps isomorphically onto the component of the moduli space of marked pairs (X, η) by forgetting E'. Consequently, the pretriangulated $K3$ category associated to the pair $(X \times X, E')$ in [MM2] depends only on the isomorphism class of X.

Contents

1. Introduction 1
1.1. Stability of modular sheaves 1
1.2. Stability in terms of the singularities along the diagonal 2
1.3. An isomorphism of two moduli spaces 3
2. Restriction to the blown-up diagonal 4
3. Proof of Theorem 1.4 8
References 10

1. Introduction

1.1. **Stability of modular sheaves.** An *irreducible holomorphic symplectic manifold* is a simply connected compact Kähler manifold, such that $H^0(X, \Omega_X^2)$ is one dimensional spanned by an everywhere non-degenerate holomorphic 2-form. Every Kähler manifold X deformation equivalent to the Hilbert scheme $S^{[n]}$ of length n subschemes of a $K3$ surface S is an irreducible holomorphic symplectic manifold [Be]. Such X is said to be of $K3^{[n]}$*-type*. Every smooth and projective moduli space $\mathcal{M}$ of stable sheaves on a $K3$ surface is an irreducible holomorphic symplectic manifold of $K3^{[n]}$-type, by results of Huybrechts, Mukai, O'Grady, and Yoshioka [O'G, Y].

Let S be a $K3$ surface, H an ample line bundle on S, v a primitive Mukai vector, and $\mathcal{M} := \mathcal{M}_H(v)$ a smooth and compact moduli space of H-stable sheaves on S of Mukai vector v. Assume that $2n := \dim_{\mathbb{C}} \mathcal{M} \geq 4$. Denote by π_S and $\pi_{\mathcal{M}}$ the two projections from $S \times \mathcal{M}$.

Date: August 23, 2016.

Let $\mathcal{U}$ be a universal sheaf over $S \times \mathcal{M}$. There is a Brauer class θ in the cohomology group $H^2_{an}(\mathcal{M}, \mathcal{O}^*_{\mathcal{M}})$, with respect to the analytic topology of $\mathcal{M}$, such that $\mathcal{U}$ is $\pi^*_{\mathcal{M}}\theta$-twisted. Let π_{ij} be the projection from $\mathcal{M} \times S \times \mathcal{M}$ onto the product of the i-th and j-th factors. Let

$$E := \mathcal{E}xt^1_{\pi_{13}}\left(\pi^*_{12}\mathcal{U}, \pi^*_{23}\mathcal{U}\right) \tag{1.1}$$

be the relative extension sheaf over $\mathcal{M} \times \mathcal{M}$. Let f_i be the projection from $\mathcal{M} \times \mathcal{M}$ onto the i-th factor. E is a reflexive $f_1^*\theta^{-1}f_2^*\theta$-twisted sheaf of rank $2n-2$, which is locally free away from the diagonal, by [M2, Prop. 4.1]. Given a Kähler class ω on $\mathcal{M}$, denote by $\tilde{\omega} := f_1^*\omega + f_2^*\omega$ the corresponding Kähler class over $\mathcal{M} \times \mathcal{M}$.

Definition 1.1. Let X be a d-dimensional compact Kähler manifold and ω a Kähler class on X. The *ω-degree* of a coherent sheaf G on X is $\deg_\omega(G) := \int_X \omega^{d-1}c_1(G)$. If G is torsion free of rank r, its *ω-slope* is $\mu_\omega(G) := \deg_\omega(G)/r$. Let E be a torsion free coherent sheaf over X, which is θ-twisted with respect to some Brauer class θ. The sheaf E is *ω-slope-semistable*, if for every subsheaf F of E, satisfying $0 < \mathrm{rank}(F) < \mathrm{rank}(E)$, we have

$$\deg_\omega\left(\mathcal{H}om(E,F)\right) \leq 0.$$

E is said to be *ω-slope-stable* if strict inequality holds above. E is said to be *ω-slope-polystable*, if it is ω-slope-semistable as well as the direct sum of ω-slope-stable sheaves.

Theorem 1.2. *The sheaf E, given in Equation (1.1), is $\tilde{\omega}$-slope-stable with respect to every Kähler class ω on $\mathcal{M}$.*

The Theorem follows from the more general Theorem 1.4 stated below. The Chern class $c_2(\mathcal{E}nd(E))$, of the sheaf E in Theorem 1.2, flatly deforms to a class of Hodge type $(2,2)$ on the cartesian square $X \times X$ of every manifold of $K3^{[n]}$-type [M2, Prop. 1.2]. This fact combines with Theorem 1.2 to imply that the sheaf E is *$\tilde{\omega}$-hyperholomorphic* in the sense of Verbitsky [V] (see also [M2, Cor. 6.11]). Verbitsky proved a very powerful deformation theoretic result for such sheaves [V, Theorem 3.19]. Associated to a Kähler class ω on $\mathcal{M}$ is a *twistor family* $\mathcal{X} \to \mathbb{P}^1_\omega$ deforming $\mathcal{M}$ [Hu1]. Verbitsky's theorem implies that E extends to a reflexive sheaf over the fiber square $\mathcal{X} \times_{\mathbb{P}^1_\omega} \mathcal{X}$ of the twistor family associated to every Kähler class ω on $\mathcal{M}$. Verbitsky's theorem, applied to the sheaf E in Theorem 1.2, is central to our proof with F. Charles of the Standard Conjectures for projective manifolds of $K3^{[n]}$-type and to our work with S. Mehrotra on pretriangulated $K3$-categories associated to manifolds of $K3^{[n]}$-type [CM, MM1, MM2].

The first example of a moduli space $\mathcal{M}$, for which the above Theorem holds, was given in [M2, Theorem 7.4]. In that example the order of the Brauer class was shown to be equal to the rank of E, and so E does not have any non-zero subsheaf of lower rank. Such a maximally twisted reflexive sheaf is thus $\tilde{\omega}$-slope-stable with respect to every Kähler class ω on X.

The special case of Theorem 1.2, where $\mathcal{M} = S^{[n]}$ is the Hilbert scheme of length n subschemes of a $K3$ surface S with a trivial Picard group, was proven earlier in [M3, Theorem 1.1(1)]. In that case it was proven that, though untwisted, E again does not have any non-zero subsheaf of lower rank.

1.2. **Stability in terms of the singularities along the diagonal.** Let X be an irreducible holomorphic symplectic manifold of complex dimension $2n > 2$. Let $\beta : Y \to X \times X$ be the

blow-up of the diagonal $\Delta \subset X \times X$. We get the diagram

$$\begin{array}{ccc} D & \xrightarrow{\iota} & Y \\ {\scriptstyle p}\downarrow & & \downarrow{\scriptstyle \beta} \\ X & \xrightarrow[\delta]{} & X \times X, \end{array}$$

where ι is the natural embedding, δ the diagonal embedding, and p the natural projection. Note that D is isomorphic to the projectivization of TX. Denote by $\ell \subset p^*TX$ the tautological line subbundle and let $\ell^\perp \subset p^*TX$ be its symplectic-orthogonal subbundle. Denote by f_i the projection from $X \times X$ onto the i-th factor. Let E be a $f_1^*\theta^{-1}f_2^*\theta$-twisted reflexive sheaf of rank $2n-2$ over $X \times X$, for some Brauer class θ on X, which is locally free away from the diagonal. Given a coherent sheaf C, denote by C_{fr} the quotient of C by its torsion subsheaf. Set $V := (\beta^*E)_{fr}(D)$.

Assumption 1.3. *Assume that V is locally free and the restriction of V to D is isomorphic to $\ell^\perp/\ell$.*

Theorem 1.4. *The sheaf E is $\tilde{\omega}$-slope-stable with respect to every Kähler class ω on X.*

The Theorem is proven in Section 3.

Remark 1.5. Assumption 1.3 holds for the sheaf E in Equation (1.1), by [M2, Prop. 4.1]. Theorem 1.2 thus follows from the more general Theorem 1.4.

1.3. **An isomorphism of two moduli spaces.** The second integral cohomology of an irreducible holomorphic symplectic manifold X comes with a non-degenerate symmetric bilinear pairing known as the *Beauville-Bogomolov-Fujiki pairing*. Its signature is $(3, b_2(X)-3)$, where $b_2(X)$ is the second Betti number [Be]. A Λ-*marking* for X is an isometry $\eta : H^2(X,\mathbb{Z}) \to \Lambda$ with a lattice Λ. The moduli space $\mathfrak{M}_\Lambda$ of isomorphism classes of Λ-marked irreducible holomorphic symplectic manifolds (X,η) is a non-Hausdorff manifold of dimension $\mathrm{rank}(\Lambda)-2$ [Hu1].

The pair $(\mathcal{M}\times\mathcal{M}, E)$ in Theorem 1.2 is known to deform to all cartesian squares $X \times X$ of manifolds of $K3^{[n]}$-type [M2, Theorem 1.3]. The sheaf E is infinitesimally rigid, $\mathrm{Ext}^1(E,E)=0$, by [MM2, Lemma 5.2]. We describe next a global analogue of these two facts.

In [MM1] we constructed a moduli space $\widetilde{\mathfrak{M}}_\Lambda$ of equivalence classes of triples (X,η,E), where X is of $K3^{[n]}$-type, $n \geq 2$, η is a Λ-marking for X, and E is a rank $2n-2$ reflexive infinitesimally rigid twisted sheaf over $X \times X$, which is $\tilde{\omega}$-slope-stable[1] with respect to some Kähler class ω on X. Two pairs (X,η,E) and (X',η',E') are *equivalent*, if there exists an isomorphism $f : X \to X'$, such that $\eta' = \eta \circ f^*$, and the sheaves $\mathcal{E}nd(E)$ and $(f\times f)^*\mathcal{E}nd(E')$ are isomorphic as sheaves of algebras. The sheaf E of every triple in $\widetilde{\mathfrak{M}}_\Lambda$ is assumed to satisfy [MM1, Condition 1.6], which implies Assumption 1.3 above. Fix a connected component $\widetilde{\mathfrak{M}}^0_\Lambda$ of $\widetilde{\mathfrak{M}}_\Lambda$ and let

$$\phi : \widetilde{\mathfrak{M}}^0_\Lambda \to \mathfrak{M}^0_\Lambda$$

be the forgetful morphism sending a triple (X,η,E) to the marked pair (X,η), where $\mathfrak{M}^0_\Lambda$ is the corresponding connected component of $\mathfrak{M}_\Lambda$. The forgetful morphism ϕ is a surjective local

[1] In [MM1] it was also assumed that the sheaf $\mathcal{E}nd(E)$ is $\tilde{\omega}$-slope-polystable, but the latter property follows from the $\tilde{\omega}$-slope-stability of E, as was proven later in [M2, Prop. 6.5] for twisted sheaves (and is well known for untwisted sheaves).

homeomorphism, which is generically injective, by [MM1, Theorems 1.9 and 6.1] (conditional on [MM1, Conj. 1.12]). Following is the main application of Theorem 1.4.

Corollary 1.6. *The above morphism ϕ is an isomorphism.*

The Corollary is proven in Section 3 (conditional on [MM1, Conj. 1.12]). Taking the quotient by the monodromy action, Corollary 1.6 may be reformulated as follows.

Corollary 1.7. *On the cartesian square $X \times X$ of every manifold X of $K3^{[n]}$-type there exists a* canonical *pair*[2] *of isomorphism classes of sheaves of algebras $\mathcal{E}nd(E)$ and $\mathcal{E}nd(E^*)$, where E is a reflexive rank $2n-2$ twisted sheaf, locally free away from the diagonal, which satisfies Assumption 1.3 and is $\tilde{\omega}$-slope-stable with respect to every Kähler class ω on X.*

The Corollary is proven in Section 3 (conditional on [MM1, Conj. 1.12]). The transposition $\tau : X \times X \to X \times X$ of the two factors pulls back $\mathcal{E}nd(E)$ to $\mathcal{E}nd(E^*)$ [MM2, Lemma 4.3]. The Corollary thus implies that the construction of [MM2], associating to a triple (X, η, E) in $\widetilde{\mathfrak{M}}^0_\Lambda$ a pretriangulated $K3$-category, results in a single equivalence class of such a category for each isomorphism class of a manifold X of $K3^{[n]}$-type with non-maximal Picard rank.

2. Restriction to the blown-up diagonal

Let E be the sheaf in Theorem 1.4 and let F be a saturated subsheaf of E of rank in the range $0 < \mathrm{rank}(F) < 2n-2$. Let $\tilde{G}$ be the image of the composition $(\beta^*F)(D) \to (\beta^*E)(D) \to V$. Denote by G the saturation of $\tilde{G}$ as a subsheaf of V. The sheaf G is reflexive, since V is locally free. The restriction of G to D maps to the restriction $\ell^\perp/\ell$ of V and we denote by G' the saturation of its image as a subsheaf of $\ell^\perp/\ell$. We provide in this section an upper bound for $\deg_{\tilde{\omega}}(\mathcal{H}om(E,F))$ in terms of the first Chern class of G' (Lemma 2.5). We then provide information on $c_1(G')$ (Lemma 2.6).

Lemma 2.1. *The equality $c_1(\mathcal{H}om(E,F)) = \beta_*(c_1(\mathcal{H}om(V,G)))$ holds.*

Proof. The higher direct image $R^i\beta_*(\mathcal{H}om(V,G))$, $i > 0$, is supported on the diagonal and so its first Chern class vanishes. The equality $c_1(\beta_*\mathcal{H}om(V,G)) = c_1(R\beta_*\mathcal{H}om(V,G))$ follows. Similarly, $c_1(\beta_*\mathcal{H}om(V,G)) = c_1(\mathcal{H}om(E,F))$, as the two sheaves coincide away from the diagonal. Hence, $c_1(\mathcal{H}om(E,F))$ is the graded summand in degree 2 of $\beta_*(ch(\mathcal{H}om(V,G))td_\beta)$, by Grothendieck-Riemann-Roch [OTT]. We have $c_1(TY) = (1-2n)[D]$ and so

$$td_\beta = td(TY)/td(T[X \times X]) = 1 + \frac{1-2n}{2}[D] + \dots$$

The statement follows, by the vanishing of $\beta_*([D])$. □

Set $h := c_1(\ell^{-1}) \in H^2(D, \mathbb{Z})$.

Lemma 2.2. $p_*(h^i) = \begin{cases} 0 & \text{if } i < 2n-1 \text{ or } i \text{ is even,} \\ 1 & \text{if } i = 2n-1, \end{cases}$

and

$$p_*(h^{2n+k}) = -c_{k+1}(TX) - \sum_{\substack{j=2 \\ j \text{ even}}}^{k-1} c_j(TX)p_*(h^{2n+k-j}), \tag{2.1}$$

[2]When $n = 2$, E^* is isomorphic to $E \otimes \det(E)^*$ and the two sheaves of algebras $\mathcal{E}nd(E)$ and $\mathcal{E}nd(E^*)$ are isomorphic.

for any positive odd integer k.

Proof. The statement is well known. The vanishing of $p_*(h^i)$, for $i < 2n-1$, follows for dimension reasons. The equality $p_*(h^{2n-1}) = 1$ is proven in [F, Appendix B.4 Lemma 9]. Consider the short exact sequence $0 \to \ell \to p^*TX \to q \to 0$ defining q. Then $c_{2n}(q) = 0$, since $\operatorname{rank}(q) = 2n-1$. On the other hand, the Chern polynomial of q satisfies

$$c_t(q) = p^*c_t(TX)/c_t(\ell) = p^*c_t(TX)(1 + h + h^2 + \cdots + h^{4n-1}).$$

Consequently, $0 = c_{2n}(q) = \sum_{j=0}^{2n} p^*c_j(TX)h^{2n-j}$ and

$$h^{2n} = -\left[\sum_{j=1}^{2n} p^*c_j(TX)h^{2n-j}\right].$$

The projection formula yields

$$p_*(h^{2n+k}) = -p_*\left(\sum_{j=1}^{2n} p^*c_j(TX)h^{2n+k-j}\right) = -c_{k+1}(TX) - \sum_{j=1}^{k} c_j(TX)p_*(h^{2n+k-j}),$$

for every positive integer k. The vanishing of $p_*(h^i)$ for even i follows, by induction, from the vanishing of $c_j(TX)$ for odd j. The recursive formula (2.1) follows. □

Lemma 2.3. *The t-th symmetric power $\operatorname{Sym}^t(TX)$ of the tangent bundle is ω-slope-stable with respect to every Kähler class ω on X, for all $t \geq 0$. The space $H^0(X, (\wedge^j TX) \otimes \operatorname{Sym}^t(T^*X))$ vanishes, for all $j \geq 0$ and all $t > 1$.*

Proof. The vector bundle TX admits a Hermite-Einstein metric whose $(1,1)$-form represents ω, for every Kähler class ω on X, by Yau's proof of the Calabi Conjecture [Hu2, Cor. 4.B.22]. In particular, TX is ω-slope-polystable with respect to every Kähler class ω. The holonomy group of the tangent bundle of an irreducible holomorphic symplectic manifold of complex dimension $2n$ is the symplectic group $Sp(n)$ [Be, Prop. 4]. Consequently, the vector bundle TX is slope-stable with respect to every Kähler class, its tensor powers are poly-stable, and the indecomposable direct summands of the tensor powers correspond to irreducible representations of $Sp(n)$. Let U be the standard representation of $Sp(n)$. The symmetric powers $\operatorname{Sym}^t(U)$ are irreducible representations, for all $t \geq 0$. If $t > 1$ then $\operatorname{Sym}^t(U)$ does not appear as a subrepresentation of the exterior product $\wedge^j U$, for any $j \geq 0$. Hence, $H^0\left(X, \left(\wedge^j TX\right) \otimes \operatorname{Sym}^t(T^*X)\right)$ vanishes, for $t > 1$. □

Lemma 2.4. *Let Z be a non-zero effective divisor on D and $[Z] \in H^2(D, \mathbb{Z})$ its class. Then*

$$\int_D p^*(\omega)^{2n-1} h^{2n-1}[Z] > 0,$$

for every Kähler class ω on X.

Proof. We have a direct sum decomposition $\operatorname{Pic}(D) = p^*\operatorname{Pic}(X) \oplus \mathbb{Z}\ell$. The restriction of $\mathcal{O}_D(Z)$ to each fiber of p is effective. Hence, $\mathcal{O}_D(Z)$ is isomorphic to $\ell^{-a} \otimes p^*L$, for some line bundle $L \in \operatorname{Pic}(X)$ and for some nonnegative integer a. The space $H^0(D, \mathcal{O}_D(Z))$ does not vanish and is isomorphic to $H^0(X, L \otimes \operatorname{Sym}^a T^*X)$. Hence, L^{-1} is a subsheaf of $\operatorname{Sym}^a T^*X$. The vector bundle $\operatorname{Sym}^a T^*X$ is an ω-slope-stable bundle with a trivial determinant, by Lemma 2.3. If $a > 0$, then L^{-1} is a subsheaf of lower rank. If $a = 0$, then L^{-1} is the ideal sheaf of a non-zero

effective divisor on X, since Z was assumed to be such. In both cases we get the inequality $\deg_\omega(L) > 0$ and so

$$\int_D p^*(\omega)^{2n-1} h^{2n-1}[Z] = \int_X \omega^{2n-1} p_*(h^{2n-1}(ah + p^*c_1(L))) = \int_X \omega^{2n-1} c_1(L) = \deg_\omega(L) > 0,$$

where the second equality is due to the vanishing of $p_*(h^{2n})$ and the equality $p_*(h^{2n-1}) = 1$ of Lemma 2.2. □

Recall that the restriction of V to D is isomorphic to $\ell^\perp/\ell$, by Assumption 1.3. Let G' be the saturation of the image in $\ell^\perp/\ell$ of the restriction to D of the subsheaf G of V. Note that the sheaves $\ell^\perp/\ell$ and G' are untwisted.

Lemma 2.5. *The following inequality holds for every Kähler class ω on X.*

$$\deg_{\tilde\omega}(\mathcal{H}om(E,F)) \leq (2n-2)\binom{4n-1}{2n}\left(\int_X \omega^{2n}\right)\int_D (p^*\omega)^{2n-1} h^{2n-1} c_1(G').$$

Proof. Let $a, b \in H^2(X,\mathbb{Z})$ be the classes satisfying $c_1(\mathcal{H}om(E,F)) = f_1^*a + f_2^*b$. We have

$$\begin{aligned}
\deg_{\tilde\omega}(\mathcal{H}om(E,F)) &= \int_{X\times X} \tilde\omega^{4n-1} c_1(\mathcal{H}om(E,F)) \\
&= \int_{X\times X} (f_1^*\omega + f_2^*\omega)^{4n-1}(f_1^*a + f_2^*b) \\
&= \binom{4n-1}{2n}\left(\int_X \omega^{2n}\right)\int_X \omega^{2n-1}(a+b) \\
&= \binom{4n-1}{2n}\left(\int_X \omega^{2n}\right)\int_X \omega^{2n-1}\delta^* c_1(\mathcal{H}om(E,F)) \\
&= \binom{4n-1}{2n}\left(\int_X \omega^{2n}\right)\int_X \omega^{2n-1}\delta^*\beta_* c_1(\mathcal{H}om(V,G)), \qquad (2.2)
\end{aligned}$$

where the last equality is by Lemma 2.1. Lemma 2.2 yields

$$\delta^*\beta_* c_1(\mathcal{H}om(V,G)) = p_*\left[h^{2n-1}p^*\delta^*(\beta_* c_1(\mathcal{H}om(V,G)))\right] = p_*\left[h^{2n-1}\iota^*\beta^*(\beta_* c_1(\mathcal{H}om(V,G)))\right].$$

The equality $\beta_*\beta^*\beta_* = \beta_*$ implies that the difference between the classes $\beta^*(\beta_* c_1(\mathcal{H}om(V,G)))$ and $c_1(\mathcal{H}om(V,G))$ belongs to the kernel of β_* and is hence a multiple of $[D]$. The vanishing $p_*(h^{2n-1}\iota^*[D]) = -p_*(h^{2n}) = 0$ yields the equality

$$\delta^*\beta_* c_1(\mathcal{H}om(V,G)) = p_*\left[h^{2n-1}\iota^*(c_1(\mathcal{H}om(V,G)))\right].$$

We have $\iota^* c_1(\mathcal{H}om(V,G)) = c_1(L\iota^*\mathcal{H}om(V,G)) = c_1([\ell^\perp/\ell]^* \otimes L\iota^*(G))$. The sheaf G is reflexive, by construction, and hence its singular locus has codimension ≥ 3 in Y. It follows that $c_1(L\iota^*G) = c_1(\iota^*G) = c_1((\iota^*G)_{fr})$. The subscheme of Y, where the rank of the homomorphism $G \to V$ is lower than the rank of G, has codimension at least 2 in Y, since G is a saturated subsheaf of V. Hence, the natural homomorphism $(\iota^*G)_{fr} \to G'$ is injective. We conclude that $c_1(L\iota^*G) + [Z] = c_1(G')$, for some effective divisor Z on D. The rank $2n-2$ vector bundle $\ell^\perp/\ell$ is symplectic, and hence $c_1(\ell^\perp/\ell) = 0$. Hence

$$\iota^* c_1(\mathcal{H}om(V,G)) = (2n-2)\left(c_1(G') - [Z]\right).$$

The Projection Formula and the two displayed formulas above yield

$$\int_X \omega^{2n-1}\delta^*\beta_* c_1(\mathcal{H}om(V,G)) = (2n-2)\int_D (p^*\omega)^{2n-1}h^{2n-1}\left(c_1(G') - [Z]\right).$$

Lemma 2.4 yields the inequality

$$\int_X \omega^{2n-1}\delta^*\beta_* c_1(\mathcal{H}om(V,G)) \leq (2n-2)\int_D (p^*\omega)^{2n-1}h^{2n-1}c_1(G').$$

Lemma 2.5 follows from the above inequality and Equation (2.2). □

Lemma 2.6. *$c_1(G') = p^*\alpha - kh$, for a non-zero class $\alpha \in H^2(X,\mathbb{Z})$ and a positive integer k.*

Proof. There exist an integer k and a class $\alpha \in H^2(X,\mathbb{Z})$ satisfying $c_1(G') = p^*\alpha - kh$, since $H^2(D,\mathbb{Z}) = p^*H^2(X,\mathbb{Z}) \oplus \mathbb{Z}h$. The integer k is positive, since the restriction of $\ell^\perp/\ell$ to each fiber of $p : D \to X$ is slope-stable, by [MM1, Lemma 7.4].

The rest of the proof is by contradiction. Assume that $\alpha = 0$. Let g be the rank of G. Then $0 < g < 2n-2$ and the top exterior power of G' yields a line subbundle of $\wedge^g[\ell^\perp/\ell]$ isomorphic to ℓ^k. In particular, the vector space

$$H^0(D, \wedge^g[\ell^\perp/\ell] \otimes \ell^{-k}) \tag{2.3}$$

contains a non-zero section.

We have the short exact sequence

$$0 \to \ell^\perp \to p^*TX \to \ell^{-1} \to 0.$$

Dualizing, we get

$$0 \to \ell \to p^*T^*X \to (\ell^\perp)^* \to 0.$$

Hence also the short exact

$$0 \to \ell \otimes \wedge^{j-1}(\ell^\perp)^* \to p^* \wedge^j T^*X \to \wedge^j(\ell^\perp)^* \to 0.$$

Dualizing the latter and tensoring by ℓ^{-t} we get the short exact sequence

$$0 \to \wedge^j(\ell^\perp) \otimes \ell^{-t} \to p^*\left(\wedge^j TX\right) \otimes \ell^{-t} \to \wedge^{j-1}(\ell^\perp) \otimes \ell^{-t-1} \to 0.$$

The inclusion of $\wedge^{j-1}(\ell^\perp) \otimes \ell^{-t-1}$ in $p^*\left(\wedge^{j-1}TX\right) \otimes \ell^{-t-1}$ yields the left exact sequence[3]

$$0 \to p_*\left[\wedge^j(\ell^\perp) \otimes \ell^{-t}\right] \to \left(\wedge^j TX\right) \otimes \mathrm{Sym}^t(T^*X) \to \left(\wedge^{j-1}TX\right) \otimes \mathrm{Sym}^{t+1}(T^*X).$$

The vector space $H^0\left(X, \left(\wedge^j TX\right) \otimes \mathrm{Sym}^t(T^*X)\right)$ vanishes, for $t > 1$, by Lemma 2.3. We conclude that $H^0\left(D, \wedge^j(\ell^\perp) \otimes \ell^{-t}\right)$ vanishes, for all pairs (j,t) of integers satisfying $j \geq 0$ and $t > 1$. The short exact sequence $0 \to \ell \to \ell^\perp \to [\ell^\perp/\ell] \to 0$ yields

$$0 \to \ell \otimes \wedge^{j-1}[\ell^\perp/\ell] \to \wedge^j\ell^\perp \to \wedge^j[\ell^\perp/\ell] \to 0.$$

Tensoring by ℓ^{-t} we get

$$0 \to \wedge^{j-1}[\ell^\perp/\ell] \otimes \ell^{1-t} \to \left(\wedge^j\ell^\perp\right) \otimes \ell^{-t} \to \wedge^j[\ell^\perp/\ell] \otimes \ell^{-t} \to 0. \tag{2.4}$$

We conclude that $H^0\left(D, \wedge^{j-1}[\ell^\perp/\ell] \otimes \ell^{1-t}\right)$ vanishes, for all pairs (j,t) of integers satisfying $j \geq 1$ and $t > 1$. The vanishing of the space (2.3) follows, by taking $j = g+1$ and $t = k+1$. This provides the desired contradiction. □

[3]Under the identification of TX with T^*X via the symplectic form, the rightmost homomorphism is the homomorphism $\left(\wedge^j TX\right) \otimes \mathrm{Sym}^t(TX) \to \left(\wedge^{j-1}TX\right) \otimes \mathrm{Sym}^{t+1}(TX)$ appearing in the Koszul complex. We will not use this fact.

3. Proof of Theorem 1.4

The proof of Theorem 1.4 reduces to that of the inequality

$$\int_D (p^*\omega)^{2n-1} h^{2n-1} c_1(G') < 0,$$

for every Kähler class ω and for every non-zero saturated proper subsheaf G' of $\ell^\perp/\ell$, by Lemma 2.5. Set $c_1(G') = p^*\alpha - kh$ as in Lemma 2.6. The left hand side of the above inequality is equal to $\int_X \omega^{2n-1}\alpha$, by the projection formula and the vanishing of $p_*(h^{2n})$.

Given a positive integer N we have the sequence of inclusions

$$p_*(G' \otimes \ell^{-N}) \subset p_*([\ell^\perp/\ell] \otimes \ell^{-N}) \subset p_*\left(\wedge^2(\ell^\perp) \otimes \ell^{-N-1}\right) \subset p_*\left(\wedge^2(p^*TX) \otimes \ell^{-N-1}\right),$$

where the second inclusion follows from the sequence (2.4). The projection formula yields the inclusion $p_*(G' \otimes \ell^{-N}) \subset \left(\wedge^2 TX\right) \otimes \mathrm{Sym}^{N+1} T^*X$. The latter is an ω-polystable vector bundle with zero first Chern class. We conclude the inequality

$$\int_X \omega^{2n-1} c_1(p_*(G' \otimes \ell^{-N})) \leq 0, \tag{3.1}$$

for every Kähler class ω on X.

The higher direct images $R^i p_*(G' \otimes \ell^{-N})$ vanish, for all N sufficiently large. Hence, there exists an integer N_0, such that

$$c_1(p_*(G' \otimes \ell^{-N})) = c_1(Rp_*(G' \otimes \ell^{-N})),$$

for all $N > N_0$. The right hand side is the graded summand $P(N)$ in $H^2(X,\mathbb{Z})[N]$ of the polynomial

$$p_*\left(ch(G')\exp(Nh)td_p\right) \in H^*(X,\mathbb{Q})[N]$$

in the variable N with coefficients in the cohomology ring. Let g be the rank of G'. We have $c_1(T_p) = 2nh$,

$$\begin{aligned} td_p &= 1 + nh + \dots, \\ ch(G') &= g + (p^*\alpha - kh) + \dots \\ exp(Nh) &= \sum_{j=0}^{4n-1} h^j \frac{N^j}{j!}. \end{aligned}$$

Hence, the graded summand of $ch(G')\exp(Nh)td_p$ in $H^{4n}(D,\mathbb{Q})$ is a polynomial of degree $2n$ in N whose first two leading terms are

$$gh^{2n}\frac{N^{2n}}{2n!} + [(p^*\alpha - kh) + gnh]h^{2n-1}\frac{N^{2n-1}}{(2n-1)!} + \dots$$

The polynomial $P(N)$ is the image of the above polynomial under p_*. The vanishing of $p_*(h^{2n})$ yields that $P(N)$ has degree $2n-1$ in N and its leading term is

$$P(N) := c_1(Rp_*(G' \otimes \ell^{-N})) = \frac{N^{2n-1}}{(2n-1)!}\alpha + \dots$$

The values of the left hand side of the inequality (3.1) for $N > N_0$ are thus the values of a polynomial in $\mathbb{R}[N]$ of degree $2n-1$, whose leading coefficient is

$$\frac{1}{(2n-1)!}\int_X \omega^{2n-1}\alpha.$$

The inequality (3.1) thus implies the inequality

$$\int_X \omega^{2n-1}\alpha \leq 0, \tag{3.2}$$

for every Kähler class ω.

Let $\mathcal{K}_X$ be the Kähler cone of X. The map $\mathcal{K}_X \to H^{2n-1,2n-1}(X,\mathbb{R})$ sending ω to ω^{2n-1} is an open map. Indeed, its differential at ω is the cup product

$$(2n-1)\omega^{2n-2}\cup : H^{1,1}(X,\mathbb{R}) \to H^{2n-1,2n-1}(X,\mathbb{R}),$$

which is an isomorphism by the Hard Lefschetz Theorem. The class α does not vanish, by Lemma 2.6. Hence, the linear functional

$$H^{2n-1,2n-1}(X,\mathbb{R}) \to \mathbb{R},$$

sending λ to $\int_X \lambda\alpha$, is an open map as well. We conclude that the image of $\mathcal{K}_X$ under the composition of these two open maps is an open subset and the inequality (3.2) is strict. This completes the proof of Theorem 1.4. □

Proof of Corollary 1.6. Any two points (X,η,E_1) and (X,η,E_2) in the fiber of ϕ are inseparable, by [MM1, Theorem 6.1]. If, furthermore, E_1 and E_2 are $\tilde{\omega}$-slope-stable, with respect to the same Kähler class ω on X, then $\mathcal{E}nd(E_1)$ and $\mathcal{E}nd(E_2)$ are isomorphic as sheaves of algebras, by [MM1, Lemma 5.3]. The sheaves E_i, $i=1,2$, are assumed to satisfy [MM1, Condition 1.6], which implies Assumption 1.3, by [MM1, Lemma 7.6]. Hence, the sheaves E_i, $i=1,2$, are $\tilde{\omega}$-slope-stable with respect to every Kähler class ω on X, by Theorem 1.4. It follows that $\mathcal{E}nd(E_1)$ and $\mathcal{E}nd(E_2)$ are isomorphic as sheaves of algebras, and the two triples (X,η,E_i), $i=1,2$, represent the same equivalence class in $\widetilde{\mathfrak{M}}^0_\Lambda$. □

Proof of Corollary 1.7. The isometry group $O(\Lambda)$ of Λ acts on $\mathfrak{M}_\Lambda$ by $g(X,\eta)\mapsto(X,g\eta)$. Let $Mon(\Lambda)\subset O(\Lambda)$ be the subgroup leaving the connected component $\mathfrak{M}^0_\Lambda$ invariant. $Mon(\Lambda)$ is determined in [M1, Theorem 1.2], which implies, in particular, that elements of $Mon(\Lambda)$ act on the discriminant group $\Lambda^*/\Lambda\cong\mathbb{Z}/(2n-2)\mathbb{Z}$ as ±1. Denote by

$$cov : Mon(\Lambda)\to\{\pm1\}$$

the corresponding character. The set of $Mon(\Lambda)$-orbits in $\mathfrak{M}^0_\Lambda$ is in bijection with the set of isomorphism classes of manifolds of $K3^{[n]}$-type. Conjugating the $Mon(\Lambda)$-action via the isomorphism ϕ of Corollary 1.6 lifts it to an action on $\widetilde{\mathfrak{M}}^0_\Lambda$. The latter action is given by $g(X,\eta,E)\mapsto(X,g\eta,E^{cov(g)})$, where $E^{cov(g)}=E$, if $cov(g)=1$, and $E^{cov(g)}=E^*$, if $cov(g)=-1$, by the proof of [MM1, Theorem 1.11]. The inverse image under ϕ of the $Mon(\Lambda)$ orbit of (X,η_0) in $\mathfrak{M}^0_\Lambda$ consists of equivalence classes of triples (X,η,E) in a unique $Mon(\Lambda)$ orbit of, say, (X,η_0,E_0), by Corollary 1.6. The above description of the monodromy action on $\widetilde{\mathfrak{M}}^0_\Lambda$ implies that $\mathcal{E}nd(E)$ is isomorphic as a sheaf of algebras to $\mathcal{E}nd(E_0)$ or $\mathcal{E}nd(E_0^*)$. □

Acknowledgements: I would like to thank Sukhendu Mehrotra and Misha Verbitsky for stimulating conversations. I thank Sukhendu Mehrotra for his careful reading of an early draft and for his helpful comments.

References

[Be] Beauville, A.: *Varietes Kähleriennes dont la premiere classe de Chern est nulle.* J. Diff. Geom. 18, p. 755–782 (1983).

[CM] Charles, F., Markman, E.: *The Standard Conjectures for holomorphic symplectic varieties deformation equivalent to Hilbert schemes of $K3$ surfaces.* Compos. Math. 149 (2013), no. 3, 481–494.

[F] Fulton, W.: *Young Tableaux.* London Math. Soc. Student Texts 35, Cambridge Univ. Press, Cambridge, 1997.

[Hu1] Huybrechts, D.: *Compact Hyperkähler Manifolds: Basic results.* Invent. Math. 135 (1999), no. 1, 63-113 and Erratum: Invent. Math. 152 (2003), no. 1, 209–212.

[Hu2] Huybrechts, D.: *Complex Geometry: An introduction.* Universitext. Springer-Verlag, Berlin, 2005.

[M1] Markman, E.: *Integral constraints on the monodromy group of the hyperkähler resolution of a symmetric product of a $K3$ surface.* Internat. J. of Math. Vol. 21, Issue: 2 (2010) pp. 169–223.

[M2] Markman, E.: *The Beauville-Bogomolov class as a characteristic class.* Electronic preprint arXiv:1105.3223v3.

[M3] Markman, E.: *Stability of a natural sheaf over the cartesian square of the Hilbert scheme of points on a $K3$ surface.* Electronic preprint arXiv:1506.06191v2.

[MM1] Markman, E., Mehrotra, S.: *A global Torelli theorem for rigid hyperholomorphic sheaves.* Electronic preprint, arXiv: 1310.5782v1.

[MM2] Markman, E., Mehrotra, S.: *Integral transforms and deformations of $K3$ surfaces.* Electronic preprint, arXiv: 1507.03108v1.

[O'G] O'Grady, K.: *The weight-two Hodge structure of moduli spaces of sheaves on a K3 surface.* J. Algebraic Geom. 6 (1997), no. 4, 599–644.

[OTT] O'Brian, N. R.; Toledo, D.; Tong, Y. L.: *Grothendieck-Riemann-Roch for complex manifolds.* Bull. Amer. Math. Soc. (N.S.) 5 (1981), no. 2, 182–184.

[V] Verbitsky, M.: *Hyperholomorphic sheaves and new examples of hyperkaehler manifolds,* alg-geom/9712012. In the book: Hyperkähler manifolds, by Kaledin, D. and Verbitsky, M., Mathematical Physics (Somerville), 12. International Press, Somerville, MA, 1999.

[Y] Yoshioka, K.: *Moduli spaces of stable sheaves on abelian surfaces.* Math. Ann. 321 (2001), no. 4, 817–884.

Department of Mathematics and Statistics, University of Massachusetts, Amherst, MA 01003, USA

E-mail address: markman@math.umass.edu